

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**




Fig. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




Fig. 6 X-ray powder pattern of Example 10




Figure 7. Differential scanning calorimetry of Example 10.




Fig. 8 Thermogravimetric Analysis (TGA) and Scanning Differential Thermal Analysis (SDTA) of Example 10




Figure 9. Apparent loss of DMF during the moisture balance measurement is reflected in the overall mass loss during the moisture absorption experiment. The salt is non-hygroscopic absorbing less than half of one percent water at 80% RH, (100.25% – 99.8% at end)




Figure 10 This sample of crystalline *S*-[2-[(1-Iminoethyl)amino]ethyl]-2-methyl-L-cysteine salicylate demonstrated no significant water absorption at 90% RH, and a

marked mass loss due to release of DMF. This sample was estimated to contain about 3.5% by weight DMF at the beginning of the experiment.




Figure 11. Isotherm plot of Example 10 of *S*-[2-[(1-Iminoethyl)amino]ethyl]-2-methyl-L-cysteine salicylate.




Fig. 12